

PATIENT INSTRUCTIONS – GASTROPANEL EXAMINATION

YOU HAVE BEEN SCHEDULED FOR GASTROPANEL EXAMINATION.

Please notice the following before the blood sample collection:

- Avoid eating, drinking and smoking at least 4 hours before sample collection.
- The study does **NOT** require discontinuation of regular medication. However, if you are taking some of the following medicines, and if a break (10 days) is not possible, you should inform about the medication and the possible break to the person taking the sample (phlebotomist) as those may affect the laboratory values.

Proton-pump inhibitors (PPI-medicines)

Esomeprazole:	Esomeprazol Orion, Esomeprazol Ratiopharm, Esomeprazol Sandoz, Nexium, Nexium 10 mg
Lansoprazole:	Gasterix, Lansoprazol Actavis 15 mg, Zolt
Omeprazole:	Losec, Losec MUPS, Omeprazol Actavis, Omeprazol Pensa, Omeprazol Ratiopharm, Omeprazol Sandoz, Omestad
Pantoprazole:	Pantoprazol Actavis, Pantoprazol Pensa, Pantoprazol Sandoz, Panzor, Somac Control, Somac
Rabeprazole:	Pariet

Data protection and data processing

Study requires processing of personal data. The processing is based on the General Data Protection Regulation (EU) 2016/679 and the National Data Protection Act (FIN) 1050/2018.

Biohit Oyj's patient register privacy policy is available on our website www.biohithealthcare.com.

Contact information:

Biohit Oyj
Laippatie 1
00880 Helsinki, Finland
Tel. +358 9 773 861
www.biohithealthcare.com