

PRODUCT NAME AND CLONE:	Monoclonal antibody to PEPSINOGEN II , Clone L10CC10
HOST:	Mouse
IMMUNOGEN:	Monoclonal antibody to pepsinogen II is derived from the hybridoma produced by fusion between myeloma cells and Balb/c spleen cells. Pepsinogen II purified from human tissue was used as an immunogen.
CLASS AND SUBCLASS:	IgG ₁
PRODUCT SPECIFICITY:	The antibody is specific to human pepsinogen II with no cross reactivity to human pepsinogen I.
PRODUCT CODE:	610056: 100 µg/ml vial 1ml 610060: 100 µg/ml vial 10ml
PRODUCT BUFFER:	PBS solution containing 1.0% (w/v) BSA and 0.09% (w/v) NaN ₃
STORAGE:	+2...+8°C
APPLICATIONS:	Immunohistochemical staining, Western blotting and enzyme-immunoassays
REFERENCES:	<ol style="list-style-type: none">1. Varis K. Surveillance of pernicious anemia. In Precancerous Lesions of the Gastrointestinal Tract. Scherlock P, Morson PC, Barbara L, Veronesi U (eds), Raven Press, New York, 189-194, 1983.2. Varis K, Samloff IM, Ihamäki T, Siurala M. An appraisal of tests for severe atrophic gastritis in relatives of patients with pernicious anemia. Dig. Dis. Sci., 24: 187-191, 1979.3. Sipponen P, Kekki M, Haapakoski J, Ihamäki T, Siurala M. Gastric cancer risk in chronic atrophic gastritis: statistical calculations of cross sectional data. Int. J. Cancer, 35: 173-177, 1985.4. Hattori Y, Tashiro H, Kawamoto T, Kodama Y. Sensitivity and specificity of mass screening for gastric cancer using the measurement of serum pepsinogens. Jpn. J. Cancer Res., 86: 1210-1215, 1995.5. Yoshihara M, Sumii K, Haruma K, Kiyohira K, Hattori N, Tanka S, Kajiyama G, Shigenobu T. The usefulness of gastric mass screening using serum pepsinogen levels compared with photofluorography. Hiroshima J. Med. Sci., 46: 81-86, 1997.6. Aoki K, Misumi J, Kimura T, Zhao W, Xie T. Evaluation of cut-off levels for screening of gastric cancer using serum pepsinogens and distribution of levels of serum pepsinogen I, II and of PG1/PGII ratios in a gastric cancer case-control study. J. Epidemiol., 7: 143-151, 1997.7. Kikuchi S, Wada O, Miki K, Nakajima T, Nishi T, Kobayashi O, Inaba Y. Serum pepsinogen as a new marker for gastric carcinoma among young adults. Research group on prevention of gastric carcinoma among young adults. Cancer, 73: 2695-2702, 1994.8. Farinati F, Di Mario F, Plebani M, Cielo R, Fanton MC, Valiante F, Masiero M, DeBoni M, Della Libera G, Burlin A. Pepsinogen A/Pepsinogen C or Pepsinogen A multiplied by gastrin in the diagnosis of gastric cancer. Ital. J. Gastroenterol., 23: 194-206, 1991.9. Borch K, Axelsson CK, Halgreen H, Damkjaer Nielsen MD, Ledin T, Szesci PB. The ratio of pepsinogen A to pepsinogen C: a sensitive test for atrophic gastritis. Scand. J. Gastroenterol., 24: 870-876, 1989.10. Yoshihara M, Sumii K, Haruma K, Kiyohira K, Hattori N, Kitadai Y, Komoto K, Tanka S, Kajiyama G. Correlation of ratio of serum papsinogen I and II with prevalence of gastric cancer and adenoma in Japanese subjects. Am. J. Gastroenterol., 93: 1090-1096, 1998.

For research use and further manufacturing only.

DS56 ver 6