Acetium®

HELPS Quit Smoking - Nicotine Free

BIOHIT HealthCare

Acetium[®] lozenge quit smoking without nicotine

Acetium[®] lozenge supports smoking cessation and can be used to alleviate tobacco dependence.

It is easy to use: Take one lozenge every time you smoke a cigarette.

Smoking cessation can be best achieved when you are motivated to quit, and it usually takes an average of 3-6 months of regular use. Acetium® lozenge is devoid of the side effects of conventional smoking intervention methods (such as nicotine dependence and possible adverse side effects of medicines).

How does Acetium[®] work?

Acetium[®] lozenge works by slowly releasing L-cysteine (a natural amino acid) into saliva. L-cysteine binds up to 90 % of cigarette smoke-derived acetaldehyde³.

The efficacy of Acetium[®] lozenge in smoking intervention has been evaluated in two clinical trials^{1,2}, with consistent results. Regular use of the lozenge during smoking increased the likelihood of smoking cessation 1.5-fold compared with placebo².

Composition

Acetium[®] lozenge only contains safe ingredients. The Acetaldehyde-binding lozenge also contains a small amount of xylitol, which reduces the acetaldehyde production by oral bacteria and promotes oral health.



Using Acetium®

Put one Acetium[®] lozenge in the mouth just before lighting a cigarette. Let the lozenge dissolve slowly in the mouth during smoking. Take another Acetium[®] lozenge if you continue smoking. The taste of cigarette eventually changes.

The effect can only be achieved when Acetium[®] lozenges are used every time you smoke and for the entire time you are exposed to tobacco smoke.



References:

1. Syrjänen K et al. Elimination of cigarette smokederived acetaldehyde in saliva by slow-release L-cysteine lozenge is an effective new method to assist smoking cessation. A randomised, double-blind, placebo-controlled intervention. Anticancer Res. 2016;36:2297-2306

2. Syrjänen K et al. Slow-release L-cystein (Acetium) lozenge is an effective new method in smoking cessation. A randomised, double-blind, placebocontrolled intervention. Anticancer Res. 2017;37:3639-3648

3. Salaspuro VJ et al. Eliminating carcinogenic acetaldehyde by cysteine from saliva during smoking. Cancer Epid Biomark Prev 2006;15:146-149

4. Secretan B et al. WHO International Agency for research on Cancer Monograph Working Group. A review of human carcinogens- Part E: tobacco, areca nut, alcohol, coal smoke, and salted fish. Lancet Oncol 10:1033-1034



Read more: www.acetium.com Order brochures: info@biohit.fi Instagram: Biohit_oyj

BIOHIT HealthCare

Innovating for Health

Acetium[®] lozenges are manufactured in Finland.